






THE UNIVERSITY OF
SYDNEY




Wave-like dark matter


Ciaran O'Hare
U. Sydney

CDMPP Annual meeting, November 2022

How light can dark matter be if it is made of fermions?



Sphere of degenerate fermions



$$v_F < v_{\text{esc}}$$

$$\Rightarrow m_{\chi} \gtrsim 100 \text{ eV}$$


“Tremaine-Gunn bound”: Pauli exclusion principle prevents you from cramming fermions lighter than $\sim 100 \text{ eV}$ into dark matter halos

Bosonic “wave-like” dark matter




Occupation number:

$$\mathcal{N} \approx (\rho_{\text{DM}}/m) \times \lambda_{\text{dB}}^3$$

$m = 1 \mu\text{eV} \longrightarrow \mathcal{N} \approx 10^{32} \longrightarrow \text{Macroscopically occupied ground state} \longrightarrow \text{Classical field}$




Wave-like dark matter mass range




So far we have not assumed anything about any couplings to the SM other than via gravity.

What can it couple to?



For more, see cajohare.github.io/AxionLimits/ → Now lists results from ~300 publications!

Schematic of a parameter space for an ultralight boson




Wave-like dark matter experiments




The axion

The “**QCD axion**” is the explanation for why the neutron doesn’t have an EDM

$d \propto$ some angle




Some angle $\rightarrow$ New field




$\rightarrow$ Tiny mass related to the energy scale f_a at which axion is created as a pseudo-Goldstone boson

$$m_a \simeq \frac{\Lambda_{\text{QCD}}^2}{f_a} \simeq 6 \text{ meV} \left(\frac{10^9 \text{ GeV}}{f_a} \right)$$

Axion-photon



Axion-fermion




All couplings suppressed by high energy scale: $g \sim f_a^{-1}$

Small couplings $\Rightarrow$ **dark matter**

Small couplings *and* small mass $\Rightarrow$ *wave-like* **dark matter**


Constraints on the existence of an axion coupled to the photon

(Nothing assumed about dark matter)




Constraints on the existence of an axion coupled to the photon

(Nothing assumed about dark matter)




Constraints on the existence of an axion coupled to the photon

(Nothing assumed about dark matter)




Searches that *do* assume axions are dark matter



Direct detection

To calculate *any* experimental signal of dark matter we need to know

1. How much dark matter there is around the Earth, ρ
2. How fast it's moving, v




For wave-like dark matter these correspond to:


Amplitude: $A = \frac{\sqrt{2\rho_a}}{m_a}$

Frequency: $\omega = m_a + \frac{1}{2}m_a v^2$
 $\approx m_a(1 + 10^{-6})$



Wave-like dark matter “lineshape”

Oscillations are coherent for 10^6 cycles, but most haloscopes observe over much longer timescales







Axion haloscopes




Axion haloscopes




RADES (CERN)




ORGAN (UWA)




QUAX (Legnaro)




CAPP (IBS)




HAYSTAC (Yale)




GrAHal (Grenoble)






Dark-SRF

2207.11346




ADMX-EFR

2203.14923




MADMAX

2003.10894



LAMPOST

2110.01582



MHz



GHz

THz

neV


μeV

meV




DM-Radio

2204.13781



ALPHA

2210.00017




BREAD


2111.12103

Dark matter new initiative (DMNI)

Snowmass '21

US putting \$ into
→ ADMX-EFR (high mass)
→ DM-Radio m^3 (low mass)






**Biggest challenge right now is
clearly that we don't know
the axion mass
→ Too much space to cover**

Sensitivity:


$$g_{a\gamma} \propto T^{-1/4}$$

However, if we knew the
correct mass, the axion could
be found tomorrow

Axion cosmology: the misalignment mechanism




Axion is the phase of a complex scalar field governed by a tilted wine-bottle potential.



Axion field rolls down from some angle and becomes a damped harmonic oscillator

$$\text{Amount of DM} \propto \theta_i^2$$





We should be able to do better than this.
But there are two big problems...

Pre-inflation: calculation is easy... but there is a free parameter θ_i
that we have no a priori knowledge of

Post-inflation: no free parameters... but things get complicated


Topological defects in the axion field in the post-inflationary scenario






Gravitational collapse of
inhomogeneities left over after
QCD phase transition leads to
the formation of $\sim \text{AU}$ sized
clumps of axions with masses
 $M \in [10^{-15}, 10^{-9}] M_{\odot}$

→ **axion miniclusters**



Eggemeier, O'Hare, Pierobon, Redondo, Wong [in prep.]




Miniclusters

“Minivoids”

Miniclusters contain $>80\%$ of the axions but make up $<1\%$ of the volume

We travel at about 0.2 mpc per year through the galaxy so are much more likely to be in the *minivoids* than the miniclusters

Implications for haloscopes



Typical density in the minivoids is ~ 0.085 of the mean density of dark matter

→ Not a nice conclusion, but it could have been much worse!


Enough axioms


Vector wave-dark matter: Dark photons

Extend SM gauge group: $SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)'$ with some gauge boson X^μ

Below EW $\rightarrow \mathcal{L} \supset -\frac{\chi}{2} F_{\mu\nu} X^{\mu\nu}$ “Kinetic mixing” with SM photon quantified by χ



Need a mass-generation mechanism, but that's it
 $\rightarrow$ Very minimal model. Can be a mediator to a dark sector coupled via millicharge, or a wave-like DM candidate in and of itself





Dark photon electrodynamics versus Axion electrodynamics

Axions source an effective current in a similar way, but only in the presence of an applied B-field, meaning DPs can be searched for with exactly the same techniques
→ **no B-field required!**



$$P_{\text{DP}} = \kappa \mathcal{G} V Q \rho_{\text{DM}} \chi^2 m_X$$

$$P_{\text{axion}} = \kappa \mathcal{G} V Q \rho_{\text{DM}} \frac{g_{a\gamma}^2 B^2}{m_a}$$


DP
limit




$$\chi m_X \leftrightarrow g_{a\gamma} B$$

Axion
limit

Green = dedicated DP searches

Red = recasted
axion searches






Enough photons

Back to axions... fermion couplings, e.g. neutron

→ Hard to beat astrophysical bounds




Why?


$$\mathcal{L} = -\frac{g_{an}}{2m_n} \partial_\mu a \underbrace{\bar{n} \gamma^5 \gamma^\mu n}_{\text{Spin-dependent}}$$

Hard to get large-enough samples of spin-polarised materials to compete with a star

Back to axions... fermion couplings, e.g. neutron
→ Hard to beat astrophysical bounds




CASPER-wind




An easier path... Scalar dark matter

$\phi(\mathbf{x}, t)$



$$\mathcal{L} = \dots + \frac{1}{4} g_\gamma \phi(\mathbf{x}, t) F_{\mu\nu} F^{\mu\nu} - g_\psi \phi(\mathbf{x}, t) \bar{\psi} \psi$$



Interaction looks like a mass term
→ e.g. time-varying electron mass


Scalar dark matter constraints

Two problems:

→ hard to beat laboratory fifth-force constraints


→ A “natural” theory with a UV cutoff at Λ_{Nat} must have:

$$g_e \lesssim \frac{4\pi m_\phi}{\Lambda_{\text{Nat}}}$$




Summary


- Wave





THE UNIVERSITY OF
SYDNEY


















Pure laboratory tests of axion-fermion fifth forces


Key point: any SM or BSM CP-violation (could be e.g. size of Jarlskog invariant of CKM matrix) could shift axion vev and generate **CP-violating** axion-fermion couplings in addition to the **CP-conserving** ones

$$\mathcal{L} \supset -a \sum_{\psi} \underline{g_p^{\psi}} (i\bar{\psi}\gamma^5\psi) - a \sum_{\psi} \underline{g_s^{\psi}} (\bar{\psi}\psi)$$




Monopole-monopole
(Spin independent)

→ e.g. tests of inverse
square law / WEP



Monopole-dipole
(Spin dependent)

→ Spin-mass forces e.g.
ARIADNE / QUAX



Dipole-dipole
(Spin dependent)


→ Forces between spin-
polarised samples

Monopole-dipole searches

Conceptually similar, spin an unpolarised source mass near to a spin-polarised target

ARIADNE


Constrains: g_p g_s (nucleon-nucleon)



QUAX

Constrains: g_p g_s (electron-nucleon)


(Latest QUAX result 2011.07100)




Challenge: stellar bounds tightly constrain g_p , and spin independent fifth force tests easily constrain g_s : so Astro x Lab bound on these coupling combos are very strong

Pure laboratory tests for monopole-dipole axion-mediated forces

Electron-nucleon coupling



Nucleon-nucleon coupling



Hard to beat the **astrophysical bounds**, but **ARIADNE** projects that it will

CP violation and the neutron electric dipole moment

Theory: Vacuum structure of QCD (instantons) generates a term

$$\mathcal{L}_{\text{QCD}} = \dots - \frac{\alpha_s}{8\pi} \theta_{\text{QCD}} G_{\mu\nu a} \tilde{G}_a^{\mu\nu}$$



Some angle, not fixed by theory,
so presumably $\mathcal{O}(1)$

Experiment: θ_{QCD} observable but bundled with phase from quark masses

$$\bar{\theta} = \theta_{\text{QCD}} + \theta_q$$

Where it sets the electric dipole moment of the neutron


$$d_n = (2.4 \pm 1.0) \bar{\theta} \times 10^{-3} e \text{ fm}$$



$$\nu_{\pm} = 2|\mu_n B \pm d_n E|$$

$$d_n$$

$$\theta_{\text{QCD}}$$



What do they see?

Recent measurement [2001.11966]

$$|d_n| < 1.8 \times 10^{-26} \text{ e cm} \quad (90\% \text{ CL})$$

$$\Rightarrow \theta \lesssim 10^{-10}$$


Conclusion: The strong interaction seems to be conserving CP when it generically shouldn't
 $\rightarrow$ **The strong CP problem**



The real problem: why are two completely unrelated numbers cancelling each other $< \text{ppb}$?


The solution, a la Peccei-Quinn

QCD vacuum energy density already has a minimum at $\theta = 0$ (Vafa-Witten theorem). However θ is just a parameter, there is no mechanism to cause it to want to minimise energy




PQ mechanism: what if there was?


The axion






- Introduce field, a , that couples to gluons $\propto (a/f_a) G\tilde{G}$. It will have a shift symmetry that can be used to cancel off any unwanted CP violation while VW theorem ensures $\langle a \rangle = 0$
- QCD vacuum implies a potential for the axion and thus a small mass

$$V(a) \approx \Lambda_{\text{QCD}}^4 \left[1 - \cos \left(\bar{\theta} + \frac{a}{f_a} \right) \right] \longrightarrow m_a \simeq \frac{\Lambda_{\text{QCD}}^2}{f_a} \simeq 6 \text{ meV} \left(\frac{10^9 \text{ GeV}}{f_a} \right)$$






Light ALP-photon oscillations in astrophysical B-fields (Galaxies, galaxy clusters, and the Milky Way)



Photon-axion mixing in a B-field

Raffelt & Stodolsky 1988
(<https://inspirehep.net/literature/253874>)

$$\mathcal{L}_{a\gamma} = \frac{1}{2} \partial_\mu a \partial^\mu a - \frac{1}{2} m_a^2 a^2 - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + g_{a\gamma} a \mathbf{E} \cdot \mathbf{B}$$



Linearised wave
equation for
photon-axion mixing

$$\left(\omega + \begin{pmatrix} \Delta_{\text{pl}} & 0 & 0 \\ 0 & \Delta_{\text{pl}} & \Delta_{\gamma a} \\ 0 & \Delta_{\gamma a} & \Delta_a \end{pmatrix} - i\partial_z \right) \begin{pmatrix} |A_\perp\rangle \\ |A_\parallel\rangle \\ |a\rangle \end{pmatrix} = 0$$

$$\Delta_{\gamma a} = \frac{g_{a\gamma} B_\perp}{2}$$


Mixing element

$$\Delta_a = -\frac{m_a^2}{2\omega}$$

Axion mass element

$$\Delta_{\text{pl}} = -\frac{\omega_{\text{pl}}^2}{2\omega}$$

Photon mass element



Axion only mixes with $A_\parallel$ so rotate into new basis and solve like neutrino oscillations:

$$\begin{pmatrix} |A'_\parallel\rangle \\ |a'\rangle \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} |A_\parallel\rangle \\ |a\rangle \end{pmatrix}$$

Probability for photon to convert to axion after travelling distance z :

$$|\langle \gamma(0) | a(L) \rangle|^2 = P_{a \rightarrow \gamma} = \sin^2(2\theta) \sin^2 \left(\frac{\Delta}{\cos(2\theta)} \right)$$

Where,

$$\theta = \frac{1}{2} \arctan \left(\frac{2\Delta_{\gamma a}}{\Delta_{\text{pl}} - \Delta_a} \right) \quad \Delta = \frac{|m_a^2 - \omega_{\text{pl}}^2|L}{4\omega}$$


Photon-axion conversion probability





For very light axions, m_a is negligible compared to ω_{pl}

$\rightarrow m_a \lesssim 10^{-12}$ eV for typical astrophysical plasmas $n_e = 10^{-3} \text{ cm}^{-3}$

$$\theta \sim g_{a\gamma} \frac{B_{\perp} \omega}{n_e}$$

$$\Delta \sim \frac{n_e L}{\omega}$$





Key issues to be resolved

Axion dark matter abundance/axion mass

- Current simulations cannot study the string scaling regime in full due to high required dynamical range, see e.g. [2007.04990]
- Not clear if the spectrum is dominated by IR modes (meaning large overproduction) or UV modes (Important for predicting the axion mass, see [2108.05368] for recent work with AMR)

Implications of an inhomogeneous axion distribution

- Formation of gravitationally bound structures “axion miniclusters”
- Do miniclusters survive to present day? Could be important for direct and indirect searches
- Can stable axion stars form inside miniclusters? Could there be signals of these e.g. fast radio bursts? May need to solve full SP system to understand this. [1804.05857]

